

PROGRESSIVE METHODS OF SURGICAL TREATMENT OF INTRACEREBRAL BRAIN TUMORS

Khusanov Zafar Toshmurodovich
Assistant, Department of Neurosurgery,
Samarkand State Medical University

Abstract

Brain tumors account for 1.8–2.3% of the structure of oncological diseases. The detection rate of primary brain tumors is 10.9–14.0 per 100 thousand population. Metastatic tumors of the brain are currently detected 5 times more often than primary tumors, and therefore metastatic tumor lesions of the brain are the most common among all tumor diseases of the central nervous system. In 50–60% of patients with brain tumors, involvement of functionally significant and vital brain structures in the pathological process is observed [1–3, 15, 17].

Introduction

The use of an integrated approach in the treatment of GM tumors ensures an increase in survival rates and an improvement in the quality of life of patients. The results of randomized studies indicate the effectiveness of surgical treatment followed by carrying out radiation therapy and chemotherapy taking into account the histobiological characteristics of tumors [4–7, 16]. Moreover, in patients after total removal, a higher therapeutic effect of adjuvant therapy is observed than in patients after partial removal of the tumor [8, 9, 19, 20]. When planning an operation and resolving questions regarding the permissible extent of tumor resection, it is necessary to determine the anatomical and topographic relationship between the tumor and functionally important zones (PVZ) of GM, the degree of their damage, possibilities restoration or compensation of impaired functions [10, 11, 18, 22, 24].

A progressive method of surgical treatment of brain tumors is operations using neuronavigation, which makes it possible to sap intracranial and intracerebral orientation with determination of the relationship of the tumor with the GM FVZ. Intraoperative use modern navigation systems allow more effectively and safely remove the tumor, achieve high quality of life indicators for operated patients [12–14, 21, 23, 25].

Purpose of the study: optimization of surgery clinical treatment of patients with tumors of the brain with using MN.

Materials and methods of research

The study included 43 patients operated on using multimodal neuronavigation (MN) for tumors of the hemispheres big brain. There were 46.6% women, men - 53.4%. The age of the patients ranged from 17 to 72 years, average 44.7 years. Functional status patients were assessed according to



clinical monitoring over time before and after surgery using the Karnofsky scale. Surgical treatment tactics were planned taking into account the data of a comprehensive clinical and neurological examination and analysis of the results of multislice computed tomography (MSCT), magnetic resonance imaging (MRI), functional magnetic resonance imaging (fMRI), magnetic resonance tractography (MR tractography), neuroimaging data intended for navigation support of intracranial surgical intervention, distributed into basic and additional. Basic data MRI and MSCT were intended to calculate stereotactic coordinates and intraoperative orientation; data from SPECT, MR venography, fMRI, and MR tractography supplemented the main images information about the localization of the tumor, functional and structural features of the anatomical formations adjacent to the lesion. MRI images in T1 and T2 modes, MSCT, MR venography, fMRI, MR tractography, SPECT if necessary combined and displayed on the station monitor in the form combined image in various combinations, depending on the information content of the received results and objectives of the study. For atraumatic and more complete removal tumors spreading into the FVZ and median structures of the brain, we used a surgical technique laser thermal destruction. Tumor areas were irradiated bordering the FVZ and conductive paths. The extent of surgical tumor resection was determined intraoperatively by comparing MN data and information obtained in real time in the form of images from a video monitor, as well as based on the results of postoperative CT and MRI

Results and Discussion

With application using multimodal neuronavigation, 43 operations to remove tumors from the cerebral hemispheres were performed. Comprehensive assessment of clinical examination data and neuroimaging results (MRI, MSCT, MR angiography, fMRI, MR tractography, SPECT) at the preoperative stage allows you to clarify the topographic characteristics of the tumor process, determine the degree of damage to the FVZ and median brain structures, get an idea about the vascularization of the tumor and its histobiological features. Using software navigation station "Stealth Station Application" Software Cranial 5" according to the main research MRI or MSCT performed spatial 3D modeling and performed stereotactic calculations. Sequential preoperative planning included segmentation and contouring of the tumor, 3D reconstruction of the tumor focus, determination of the area of perifocal spread edema, construction of volumetric topographic images of the surface relief of the cerebral hemispheres, convexitally located vessels, lateral ventricles, as well as the choice of the optimal trajectory and boundaries of surgical access. System The virtual image on the navigation station monitor was supplemented with video monitoring data in real time [18]. Subtotal tumor removal was performed in 252 (56.9%) patients, total - in 161 (36.3%), partial - in 30 (6.8%). According to the results of histological examination of intraoperative biopsy material, glioma grades I and II anaplasia (WHO) detected in 71 (16.0%) patients, anaplastic glioma grade III anaplasia - in 138 (31.2%), glioma IV degree of anaplasia - in 167 (37.7%), metastatic tumors - in 67 (15.1%). After surgical treatment, the assessment of the patients' functional status increased on average from 68.4 to 86.2 points on the Karnofsky scale. Depending on the location of the tumors according to in relation to the FVZ of the cerebral hemispheres, in according to topographic classification R. Sawaya [19], three groups of tumors were identified: with lesions of the FVZ (Sawaya Grade III) - in 55.5% patients adjacent to the



FVZ (Sawaya Grade II) – in 35.2%, located remotely from the FVZ (Sawaya Grade I) - 9.3%. Total removal of Sawaya Grade I tumors produced in 26.2% of patients, Sawaya Grade III - in 42.7% %; subtotal - in 67.3 and respectively 49.6%; partial - in 6.5 and 7.7%. After surgery functional status of patients in both groups almost identical, in the Sawaya Grade II group - 88.9 points, Sawaya Grade III - 85.6 points. Strategy for surgical treatment of patients with regarding tumors of the cerebral hemispheres suggests optimizing the volume of tumor resection with ensuring high survival rates patients while maintaining high quality life. It is fundamentally important to strive for maximum tumor removal and reducing the risk of possible postoperative neurological deficit [4, 10, 20, 21].

Conclusions

1. Comprehensive analysis of MN data on stages of preoperative planning allows determine anatomical and topographical features tumor spread, get information about the nature and extent of damage to the FVZ and median structures of the GM, to identify changes in structures GM due to growth and volumetric action tumors. 2. Preoperative multimodal navigation planning using integrated neuroimaging data and modeling in virtual computer space of anatomical and functional formations allows you to select trajectory and safe limits of surgical access, determine the optimal volume of resection tumor tissue, as well as assess the risk of possible intraoperative trauma to the cerebral hemorrhage. 3. Comparison of virtual preoperative navigation planning data with real topographic relationships of anatomical structures in the surgical field provides intraoperative orientation and controlled performing surgical procedures with simultaneous recording of the stages of the operation using video monitoring systems. 4. Intraoperative support using MN ensures high precision of surgical manipulations and functional preservation of the structures of the brain, which helps to reduce degree of risk of postoperative neurological deficit.

References

1. Rozumenko V.D. Neuro-oncology: current status problems / V.D. Rozumenko // *Oncology*. - 2006. - T.8, No. 2. — P.188–191.
2. Hoffman S. Temporal trends in incidence of primary brain tumors in the United States, 1985–1999 / S. Hoffman, M. Jennifer, J. Bridget // *NeuroOncology*. - 2006. - V.8, N1. — P.27–37.
3. Intraoperative subcortical language tract mapping guides surgical removal of gliomas involving speech areas / L. Bello, M. Gallucci, M. Fava, G. Carrabba, C. Giussani, F. Acerbi, P. Baratta, V. Songa, V. Conte, V. Branca, N. Stocchetti, C. Papagno, S.M. Gaini // *Neurosurgery*. - 2007. - V.60, N1. — P.67–80.
4. Motor functional MRI for preoperative and intraoperative neurosurgical guidance / ID Wilkinson, CA Romanowski, D. A. Jellinek, J. Morris, P. D. Griffiths // *Br. J. Radiol*. — 2003. - V.76, N902. — P.98–103.
5. Treatment outcomes and prognostic factors in patients with supratentorial low-grade gliomas / SA Ho, JT Lui, CC Huang, YJ Huang, CY Hsiung, EY Huang // *Br. J. Radiol*.



- 2005. - V.78, N927. — P.230–235.

6. Barker FG Surgery for primary supratentorial brain tumors in the United States, 1988 to 2000: the effect of provider caseload and centralization of care / FG Barker, WT Curry, BS Carter // *NeuroOncology* - 2005. - V.7, N1.

— P.49–63.

7. Present day's standards in microsurgery of low-grade gliomas / L. Bello, E. Fava, G. Carrabba [et al.] // *Advances and Technical Standards in Neurosurgery*. Vol.35. Low-Grade Gliomas; ed. J. Schramm. — Wien: Springer-Verlag, 2010.

— P.113–158.

8. A multivariate analysis of 416 patients with glioblastoma multiforme: Prognosis, extent of resection, and survival / M. Lacroix, D. Abi-Said, D. R. Fournay, Z. L. Gokaslan, W. Shi, F. DeMonte, F. F. Lang, I. E. McCutcheon, S. J. Hassenbusch,

E. Holland, K. Hess, C. Michael, D. Miller, R. Sawaya // *J. Neurosurg.* - 2001. - V.95, N2. — P.190–198.

9. Extent of resection and survival in glioblastoma multiforme:

Identification of and adjustment for bias / W. Stummer, H. J. Reulen, T. Meinel, U. Pichlmeier, W. Schumacher, J. C. Tonn, V. Rohde, F. Oppel, B. Turowski, C. Woiciechowsky, K. Franz, T. Pietsch // *Neurosurgery*. - 2008. - V.62, N3.

— P.564–576.

10. Gil-Robles S. Surgical management of World Health Organization Grade II gliomas in eloquent areas: the necessity of preserving a margin around functional structures / S. Gil-Robles, H. Duffau // *Neurosurg. Focus*. — 2010.

— V.28, N2. - E.8.

11. González-Darder JM Management of brain tumors in eloquent areas / J. M González-Darder, P. González-López // *Diagnostic techniques and surgical management of brain tumors*; ed. AL Abujamra. — Croatia: InTech, 2011.

— P.455–492.

12. Effectiveness of neuronavigation in resecting solitary intracerebral contrast-enhancing tumors: a randomized controlled trial / PW Willems, MJ Taphoorn, H. Burger, JW Berkelbach van der Sprenkel, CA Tulleken // *J. Neurosurg.*

- 2006. - V.104, N3. — P.360–368.

13. Image-guided resection of high-grade glioma: Patient selection factors and outcome / N. Scottlitofsky, A. M. Bauer, RS Kasper, CM Sullivan, OH Dabbous // *Neurosurg. Focus*.

— 2006.— V.20, N4. - E.16.

14. Impact of neuronavigation and image-guided extensive resection for adult patients with supratentorial malignant astrocytomas: a single-institution retrospective study / M.

15. Tychiev L. N. et al. NASOPHARYNGEAL EXTRACTION OF S. PNEUMONIAE FROM ADULT PATIENTS WITH ACUTE RESPIRATORY INFECTIONS AND ANTIBIOTIC RESISTANCE OF ISOLATED STRAINS // *Art of Medicine. International Medical Scientific Journal*. – 2022. – T. 2. – №. 1.



16. Раббимова Н. Т., Матякубова Ф. Э., Тиркашев О. С. ЧАСТОТА ВЫДЕЛЕНИЯ STREPTOCOCCUS PNEUMONIAE ПРИ ОСТРЫХ РЕСПИРАТОРНЫХ ИНФЕКЦИЯХ ДЫХАТЕЛЬНЫХ ПУТЕЙ //VOLGAMEDSCIENCE. – 2021. – С. 589-591.
17. Tuychiev L. N. et al. Antimicrobial susceptibility OF S. Pneumoniae, isolated from adults //湖南大学学报 (自然科学版). – 2021. – Т. 48. – №. 11.
18. Раббимова Н. и др. Математическое моделирование и прогнозирование заболеваемости кожным лейшманиозом в республике узбекистан //Журнал проблемы биологии и медицины. – 2017. – №. 1 (93). – С. 104-107.
19. Сувонкулов У. и др. Идентификация видовой принадлежности возбудителей кожного лейшманиоза методом полимеразной цепной реакции //Журнал проблемы биологии и медицины. – 2016. – №. 3 (89). – С. 91-92.
20. Egamovna M. F. et al. CLINICAL AND EPIDEMIOLOGICAL FEATURES OF THE COURSE OF SHIGELLOSIS IN ADULTS AT THE PRESENT STAGE IN 2009-2019 //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 5. – С. 1285-1294.
21. Абдухалилова Г. К. и др. Назофарингеальное носительство str. e у взрослых. – 2022.
22. Egamovna M. F. et al. CLINICAL AND EPIDEMIOLOGICAL FEATURES OF THE COURSE OF SHIGELLOSIS IN ADULTS AT THE PRESENT STAGE IN 2009-2019 //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 5. – С. 1285-1294.
23. Абдухалилова Г. К. и др. Динамика устойчивости к антибиотикам и частота назофарингеального выделения S. Pneumoniae у взрослых с острыми респираторными инфекциями. – 2022.
24. Ярмухамедова Н. и др. Особенности течения хронического гепатита с на фоне туберкулеза //Журнал вестник врача. – 2019. – Т. 1. – №. 1. – С. 129-132.1
25. Anvarovna, Y. N., Egamovna, M. F., Tashtemirovna, R. N., Buribayevna, M. G., & Saidovich, T. O. (2021). Clinical and Epidemiological Characteristics of Shigellosis in Adults at the Contemporary Stage. Central Asian Journal of Medical and Natural Science, 2(3), 311-318. <https://doi.org/10.47494/cajms.v2i3.221>
26. Тиркашев, О. С. Клинико-эпидемиологическая характеристика кори в Самаркандской области / О. С. Тиркашев, Ф. Э. Матякубова, Н. Т. Раббимова // VOLGAMEDSCIENCE : Сборник тезисов VII Всероссийской конференции молодых ученых и студентов с международным участием: материалы конференции, Нижний Новгород, 16–18 марта 2021 года. – Нижний Новгород: Федеральное государственное бюджетное образовательное учреждение высшего образования "Приволжский исследовательский медицинский университет" Министерства здравоохранения Российской Федерации, 2021. – С. 624-625. – EDN GZYHJQ.

