

COVID-19 MEDICAL REHABILITATION FOR PATIENTS WITH COVID-19

Abdurahimova Zulfiya Qudratullayevna
Nilufar Kurvankulovna Djurayeva

Abstract

The COVID-19 pandemic has posed new and complex challenges to healthcare systems around the world. This article examines the organization of medical rehabilitation for patients infected with COVID-19, the stages of the rehabilitation process, existing problems, and solutions to overcome them. The goal is to improve the quality of life of patients and restore their ability to work through effective medical rehabilitation.

Keywords: COVID-19, rehabilitation, pandemic, medical staff.


Introduction

Medical rehabilitation is essential for full recovery in patients with coronavirus pneumonia. Rehabilitation should begin after the patient's condition stabilizes. It is initially continued in intensive care units, then in hospitals, at home, and in sanatorium-resort institutions.

Organization of the rehabilitation process

A multidisciplinary rehabilitation team (MDR) should be formed to provide rehabilitation services to patients with COVID-19. This team should include:


- Attending physician,
- Physiotherapist,
- Physical therapy doctor,
- Instructor-methodologist,
- Ward nurse and other specialists.



Problems and solutions

- The isolation period during the recovery period (14 days) can delay rehabilitation.
- Lack of specialists involved in rehabilitation.
- Mental and physical fatigue of medical workers working in the “red zone”.

COVID-19 Rehabilitation Algorithm for Patients



Solutions:


- Organization of training via telemedicine.
- Acceptance of rehabilitation as an integral part of treatment.
- Scheduling of work activities and psychological support for medical workers.

Scientific and practical results

During the study, a 3-stage rehabilitation system was introduced, and active recovery was achieved in patients with COVID-19, who also had problems with the central and peripheral nervous system, cardiovascular system, and musculoskeletal system. A multidisciplinary approach served to increase efficiency.



Stages of rehabilitation



Conclusion

COVID-19 has become a serious test for the world's healthcare system. A complete cure for this disease requires not only getting rid of the virus, but also complete rehabilitation. Through targeted and systematic rehabilitation measures, it is possible to improve the quality of life of patients, return them to a healthy lifestyle, and restore their ability to work. Therefore, specialists must develop a medical rehabilitation algorithm for patients infected with COVID-19.

REFERENCES

1. Wang S., Pan R., Wang H., Tang Y., Xu L., Ho X.S. and etc . Immediate psychological reactions and associated factors during the initial stage of the coronavirus disease (COVID-19) epidemic among the Chinese population in 2019. *Int J Environ Res Public Health*. 2020; 17 (5). Electronic publication 03/12/2020. doi : 10.3390/ijerph17051729; PubMed Central PMCID: PMC7084952. [Free article PMC] [PubMed] [CrossRef] [Google Scholar]
2. Ahmed MZ, Ahmed O, Aibao Z, et al. Epidemic of covid-19 in China and associated psychological problems. // *Asian J Psychiatr* . 2020;51:102092.
3. Arasli, H., Furunes, T., Jafari, K., Saydam, M.B., Degirmencioglu, Z. Hearing the voices of wingless angels: A critical content analysis of nurses' Covid-19 experiences. // *International Journal of Environmental Research and Public Health*. 2020; 17(22); 8484: 1-16.





4. Atay , S., Cura , ŞÜ. Problems Encountered by Nurses Due to the Use of Personal Protective The current document contains no sources. 2020; 66(10): 12-16.
5. Bostan, S., Akbolat, M., Kaya, A., Ozata, M., Gunes, D. Assessments of anxiety levels and working conditions of health employees working in COVID-19 pandemic hospitals //Electronic Journal of General Medicine. 2020; 17(5): em246.
6. Brugliera L, Spina A, Castellazzi P, et al. Rehabilitation of Covid-19 patients. //J Rehabil Med. 2020; 52(4): jrm00046.
7. Study of factors influencing the effectiveness of rehabilitation during COVID-19. authors Tuychiev L.N., Khudaykulova G.K., Eraliev U.E., Dzhuraeva N.K., Sadullaev S.E.
8. Negrini S, Ferriero G, Kiekens C, Boldrini P. Facing in real time the challenges of the Covid-19 epidemic for rehabilitation. Eur J Phys Rehabil Med. 2020. doi : 10.23736/S1973-9087.20.06286-3.
9. The role of rehabilitation in primary health care Authors Tuychiev Laziz N., Eraliev Umid E., Khudaykulova Gulnara K., Juraeva Nilufar
10. Simpson R, Robinson L. Rehabilitation following critical illness in people with COVID-19 infection. Am J Phys. Med Rehabil . 2020;99(6):470–474. doi 10.1097/PHM.0000000000001443.
11. Psychological component of medical rehabilitation of patients with the post-Covid syndrome, Juraeva N.K., conference, 23–29
12. LN.Tuychiev, GK Khudaikulova, N Djuraeva - 2023
13. Rol Reabilitatsii V Usloviyakh Pandemii . U. Eraliev, G. Khudaikulova, A. Usmankhodzhaev - Evraziysky zurnal meditsinskih i stroitelnyx ..., 2022
14. I am improving the performance of nurses in the rehabilitation of patients after E Eraliev, N.K Djuraeva
15. Rust G, Melbourne M, Truman BI, Daniels E, Fry-Johnson Y, Curtin T. Role of the primary care safety net in pandemic influenza. Am J Public Health. 2009;99

