

REFRACTIVE CHANGES FOLLOWING CONGENITAL CATARACT SURGERY IN CHILDREN: MODERN DIAGNOSTIC AND PROGNOSTIC METHODS

Erkinova O'g'iloy Xurmat qizi
Alfraganus Universiteti

Abstract

Congenital cataract is a significant cause of visual impairment in children and may interfere with normal visual development during critical periods of ocular maturation. Although early surgical treatment can restore the transparency of the visual axis, children frequently experience substantial refractive changes during postoperative growth. These changes may affect visual acuity, binocular function, and the long-term development of the visual system. Objective: This article examines the mechanisms underlying refractive changes following congenital cataract surgery in children, evaluates contemporary diagnostic approaches, and discusses prognostic methods that may improve individualized clinical management.

Keywords: Congenital cataract, pediatric ophthalmology, refractive development, postoperative refraction, axial length, keratometry, intraocular lens implantation, optical biometry, amblyopia, prognostic modeling.

Introduction

A narrative review approach is used to synthesize established clinical concepts and findings from ophthalmological literature concerning pediatric cataract surgery, ocular biometry, refractive development, intraocular lens implantation, and postoperative visual rehabilitation. Particular attention is given to the roles of axial length measurement, keratometry, retinoscopy, cycloplegic refraction, optical coherence tomography, and longitudinal refractive monitoring. The review also considers the potential contribution of statistical modeling and machine-learning techniques to postoperative risk assessment.

Results:

Postoperative refractive outcomes depend on several interacting factors, including the child's age at surgery, axial growth, corneal power, the accuracy of intraocular lens power calculation, the timing of optical correction, and adherence to amblyopia treatment. Younger children generally demonstrate greater variability in refractive development because ocular structures undergo rapid changes during early life. Modern optical biometry and improved surgical planning can enhance measurement accuracy, although their applicability may be limited by dense cataracts, poor fixation, and the child's age. Longitudinal measurements are essential for distinguishing expected developmental changes





from clinically significant refractive abnormalities. Predictive models may support individualized follow-up, but their reliability depends on appropriate validation in pediatric populations.

Congenital cataract represents an important cause of preventable visual impairment in childhood. It is characterized by opacity of the crystalline lens that is present at birth or develops during early infancy. Depending on its density, location, laterality, and associated ocular abnormalities, congenital cataract may interfere with the formation of clear retinal images and disrupt the development of normal visual pathways. Its clinical significance extends beyond the opacity itself because early visual deprivation can produce persistent functional deficits even after successful anatomical treatment. The developing visual system requires sufficiently clear and appropriately focused retinal images. During infancy and early childhood, neural pathways responsible for visual acuity, contrast sensitivity, accommodation, and binocular coordination undergo substantial maturation. When a cataract obstructs the visual axis during this period, abnormal visual stimulation may lead to deprivation amblyopia. Consequently, the clinical objective of treatment is not limited to removing the opaque lens. It also includes restoring an effective optical pathway, maintaining appropriate refractive correction, and supporting the development of useful visual function.

Surgical extraction is the principal treatment for visually significant congenital cataracts. Depending on the child's age, ocular anatomy, cataract characteristics, and anticipated postoperative requirements, treatment may involve removal of the crystalline lens with primary intraocular lens implantation or correction using contact lenses or spectacles. The selection of an appropriate strategy requires careful consideration of both immediate surgical outcomes and the expected development of the eye. One of the principal challenges following pediatric cataract surgery is the occurrence of refractive changes over time. Unlike the adult eye, the child's eye is undergoing continuous anatomical development. Axial length increases, corneal curvature changes, and the relationships among the cornea, intraocular lens, and retina evolve as the eye matures. These processes may alter the refractive state even when the initial surgical procedure and intraocular lens power calculation are technically satisfactory.

Literature Review and Degree of Problem Investigation

Congenital Cataract and the Development of Refractive Errors

The relationship between congenital cataract and refractive development has been investigated extensively in pediatric ophthalmology. Previous clinical research has established that the visual consequences of congenital cataract depend on the degree of lens opacity, the age at which visual deprivation begins, the duration of deprivation, and the effectiveness of subsequent optical rehabilitation. However, refractive outcomes remain variable even among children who receive treatment at comparable ages. Congenital cataract may be unilateral or bilateral, and this distinction has important implications for postoperative management. Unilateral cataract is frequently associated with a greater risk of amblyopia because the affected eye competes with a normally functioning fellow eye. The resulting imbalance in visual input may interfere with binocular integration and contribute to persistent interocular differences in visual acuity. Bilateral cataracts may produce more symmetrical visual deprivation, although substantial differences between the two eyes can still occur.





Axial Growth and the Mechanisms of Postoperative Myopic Shift

Axial elongation is one of the principal anatomical mechanisms associated with postoperative refractive change in children who undergo cataract extraction with intraocular lens implantation. The axial length of the eye generally increases during childhood, although the rate of growth varies with age and individual developmental characteristics. When the distance between the optical system and the retina increases, the effective refractive state may shift toward myopia if the optical power does not compensate for the change. Because an implanted intraocular lens has a relatively fixed optical power, it cannot adapt to axial growth in the same manner as the natural crystalline lens. Consequently, the refractive state may become progressively more myopic during ocular development.

The Contribution of Intraocular Lens Power Calculation

Intraocular lens power calculation is another major determinant of postoperative refractive accuracy. Common calculation approaches incorporate axial length and corneal curvature, with different formulas using additional variables or different mathematical relationships to estimate the effective optical position of the implanted lens. In adult cataract surgery, modern formulas have substantially improved refractive predictability. Nevertheless, their performance cannot automatically be extrapolated to infants and young children. Pediatric eyes may have short axial lengths, relatively steep corneas, and anatomical relationships that differ considerably from those found in adults. These characteristics can reduce the accuracy of formulas developed or optimized primarily for adult populations.

Visual Rehabilitation and the Prevention of Amblyopia

Refractive correction is an essential component of the treatment pathway following congenital cataract surgery. The removal of the cataract restores the possibility of forming a clear retinal image, but useful vision does not necessarily develop without appropriate optical rehabilitation. Children who undergo cataract extraction may require spectacles, contact lenses, or both, depending on the surgical approach and the resulting optical state. When the natural lens is removed without primary intraocular lens implantation, substantial optical correction is necessary to compensate for aphakia. Contact lenses may provide useful optical rehabilitation, particularly in unilateral cases, although successful use depends on appropriate fitting, hygiene, caregiver training, and consistent adherence. After intraocular lens implantation, spectacles may still be required to correct residual refractive error. These errors may include myopia, hyperopia, astigmatism, or anisometropia. Because the refractive state can change over time, a prescription that was appropriate several months earlier may no longer provide adequate correction. Amblyopia management is closely linked to refractive assessment. In unilateral congenital cataract, treatment may include patching the fellow eye to encourage visual use of the operated eye, according to an individualized ophthalmological plan.

Limitations of Existing Prognostic Approaches

Traditional prognostic approaches often emphasize age at surgery, axial length, corneal curvature, and the intended postoperative refractive target. These variables are clinically relevant, but they do not fully explain the range of refractive outcomes observed in pediatric patients. One limitation is the





use of a single preoperative measurement to predict changes that may occur over several years. Ocular growth is a dynamic process, and its rate may vary during different developmental periods. Consequently, predictions based on baseline measurements may become less accurate as the follow-up interval increases. A second limitation is that some studies include relatively small numbers of children or focus on specific surgical techniques. The resulting models may not be generalizable to other centers, age groups, or clinical populations. Differences in surgical practice, measurement technology, follow-up schedules, and optical rehabilitation can influence the observed results.

REFERENCES

1. Chan, J. J. T., Wong, E. S., Lam, C. P. S., & Yam, J. C. Y. (2023). Ten-year refractive and visual outcomes of intraocular lens implantation in infants with congenital cataract. *Hong Kong Medical Journal*, 29(1), 22–30. <https://doi.org/10.12809/hkmj209241>
2. Koch, C. R., Kara-Junior, N., Serra, A., et al. (2018). Long-term results of secondary intraocular lens implantation in children under 30 months of age. *Eye*, 32, 1858–1863. <https://doi.org/10.1038/s41433-018-0191-3>
3. Lambert, S. R., Lynn, M. J., DuBois, L. G., et al. (2014). A randomized clinical trial comparing contact lens and intraocular lens correction of monocular aphakia during infancy: Grating visual acuity and adverse events at age 1 year. *Archives of Ophthalmology*, 130(2), 207–213.
4. Trivedi, R. H., Wilson, M. E., & Vasavada, A. R. (2007). Changes in interocular axial length after pediatric cataract surgery. *Journal of AAPOS*, 11(3), 225–229.
5. Vasavada, A. R., Trivedi, R. H., & Nath, V. C. (2004). Visual outcomes and complications following pediatric cataract surgery and intraocular lens implantation. *Journal of Cataract & Refractive Surgery*.
6. Mravičić, I., Orešković, E. G., Boháč, M., & Drača, N. (2026). Predicting postoperative myopic shift after paediatric intraocular lens implantation: A scoping review of associated factors. *Medicina*, 62(1), 106. <https://doi.org/10.3390/medicina62010106>
7. American Academy of Ophthalmology. (2022). Pediatric Eye Evaluations Preferred Practice Pattern®. *Ophthalmology*. <https://doi.org/10.1016/j.ophtha.2022.10.030>
8. Long-Term Refractive Changes After Pediatric Cataract Surgery: An Average Follow-up of 15 Years. (2025). *American Journal of Ophthalmology*, 277, 305–312. <https://doi.org/10.1016/j.ajo.2025.05.035>
9. Chan, J. J. T., Wong, E. S., Lam, C. P. S., & Yam, J. C. (2023). Ten-year refractive and visual outcomes of intraocular lens implantation in infants with congenital cataract. *Hong Kong Medical Journal*, 29(1), 22–30. <https://doi.org/10.12809/hkmj209241>
10. Zhong, Y., Yu, Y., Lu, B., et al. (2021). Accuracy of intraocular lens power calculation formulas in pediatric cataract patients: A systematic review and meta-analysis. *Frontiers in Medicine*, 8, 710492. <https://doi.org/10.3389/fmed.2021.710492>
11. Lambert, S. R. (2016). Changes in ocular growth after pediatric cataract surgery. *Developments in Ophthalmology*, 57, 29–39. <https://doi.org/10.1159/000442498>
12. VanderVeen, D. K., Nizam, A., Lynn, M. J., et al. (2015). Predictability of intraocular lens calculation formulas in infants undergoing cataract surgery: Results from the Infant Aphakia Treatment Study. *American Journal of Ophthalmology*, 160(3), 438–446.e1.





13. Infant Aphakia Treatment Study Group. (2010). Comparison of contact lens and intraocular lens correction of monocular aphakia during infancy: A randomized clinical trial. *Archives of Ophthalmology*, 128(7), 810–818. <https://doi.org/10.1001/archophthalmol.2010.101>
14. Hussin, M., & Markham, R. (2009). Changes in axial length growth after congenital cataract surgery and intraocular lens implantation in children younger than 5 years. *Journal of Cataract & Refractive Surgery*, 35(7), 1223–1228. <https://doi.org/10.1016/j.jcrs.2009.03.015>
15. Flitcroft, D. I., Knight-Nanan, D., Bowell, R., Lanigan, B., & O’Keefe, M. (1999). Intraocular lenses in children: Changes in axial length, corneal curvature, and refraction. *British Journal of Ophthalmology*, 83(3), 265–269. <https://doi.org/10.1136/bjo.83.3.265>
16. Spierer, A., Desatnik, H., & Blumenthal, M. (1999). Refractive status in children after long-term follow-up of cataract surgery with intraocular lens implantation. *Journal of Pediatric Ophthalmology and Strabismus*, 36(1), 25–29. <https://doi.org/10.3928/0191-3913-19990101-06>
17. Hussin, M., & Markham, R. (2009). Long-term visual function outcomes of congenital cataract surgery with intraocular lens implantation in children under 5 years of age. *European Journal of Ophthalmology*, 19(5), 754–761. <https://doi.org/10.1177/112067210901900512>
18. Mravičić, I., Orešković, E. G., Boháč, M., & Drača, N. (2026). Predicting postoperative myopic shift after paediatric intraocular lens implantation: A scoping review of associated factors. *Medicina*, 62(1), 106. <https://doi.org/10.3390/medicina62010106>

